

# Dissecting the pleiotropic consequences of a quantitative trait nucleotide

Hyun Seok Kim, Juyoung Huh &amp; Justin C. Fay

Department of Genetics, Washington University School of Medicine, St. Louis, MO, USA

**Correspondence:** Justin C. Fay, Department of Genetics, Washington University School of Medicine, 444 Forest Park Ave, St. Louis, MO 63108, USA. Tel.: +1 314 747 1808; fax: +1 314 362 2156; e-mail: jfay@genetics.wustl.edu

**Present address:** Hyun Seok Kim, Department of Cell Biology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

Received 13 January 2009; revised 23 March 2009; accepted 24 March 2009.  
Final version published online 23 April 2009.

DOI:10.1111/j.1567-1364.2009.00516.x

Editor: Ian Dawes

## Keywords

QTN; *CYS4*; pleiotropy; gene expression; hydrogen sulfide.

## Introduction

The molecular basis of phenotypic variation in natural populations is important to understand both molecular and phenotypic evolution. Yet, dissecting the molecular basis of a trait is difficult as it involves not only mapping quantitative trait polymorphisms but also understanding how each polymorphism affects a trait by influencing the activity of other genes, pathways and cellular processes involved in the production of a trait. For example, a quantitative trait polymorphism in *MKT1* affects high-temperature growth (Sinha *et al.*, 2006), sporulation efficiency (Deutschbauer & Davis, 2005), DNA damage sensitivity (Demogines *et al.*, 2008), as well as numerous gene expression levels (Smith & Kruglyak, 2008), but the mechanisms by which the *MKT1* polymorphism affects these traits have only begun to be characterized (Lee *et al.*, 2009).

The relationship among molecular, cellular, and organismal changes caused by quantitative trait polymorphisms is

## Abstract

The downstream consequences of a single quantitative trait polymorphism can provide important insight into the molecular basis of a trait. However, the molecular consequences of a polymorphism may be complex and only a subset of these may influence the trait of interest. In natural isolates of *Saccharomyces cerevisiae*, a nonsynonymous polymorphism in cystathione  $\beta$ -synthase (*CYS4*) causes a deficiency in both cysteine and glutathione that results in rust-colored colonies and drug-dependent growth defects. Using a single-nucleotide allele replacement, we characterized the effects of this polymorphism on gene expression levels across the genome. To determine whether any of the differentially expressed genes are necessary for the production of rust-colored colonies, we screened the yeast deletion collection for genes that enhance or suppress rust coloration. We found that genes in the sulfur assimilation pathway are required for the production of rust color but not the drug-sensitivity phenotype. Our results show that a single quantitative trait polymorphism can generate a complex set of downstream changes, providing a molecular basis for pleiotropy.

key to obtaining a general understanding of the molecular basis of a trait. Changes in gene expression provide a rich measurement of molecular variation and have provided valuable insights into the genes and molecular processes related to a trait of interest, for example (Cavaliere *et al.*, 2000; de Haan *et al.*, 2002; Schadt *et al.*, 2003; Fay *et al.*, 2004). However, the relationship between gene expression and other phenotypes may often be complex; changes in gene expression that are tightly correlated with a trait may not be involved in the production of the trait. In yeast, changes in the environment cause many genes to change expression even though most are not required to survive or grow in the new environment (Birrell *et al.*, 2002; Giaever *et al.*, 2002). However, environmentally induced changes in gene expression may be different from genetic variation in gene expression segregating in natural populations because the latter is heritable and shaped by natural selection. Thus, while there are numerous examples of changes in gene expression with phenotypic consequences (Carroll *et al.*,

2001; Wray *et al.*, 2003), relatively little is known about the extent to which the effects of a quantitative trait polymorphism are mediated by downstream changes in gene expression.

Identifying the molecular consequences of a single quantitative trait polymorphism is difficult when there are other polymorphisms segregating within the genetic background. In yeast, allele-replacement strains have been used to measure the precise effects of a number of quantitative trait polymorphisms on gene expression levels (Yvert *et al.*, 2003; Brown *et al.*, 2008; Smith & Kruglyak, 2008). Each polymorphism resulted in numerous gene expression changes, but the role that these changes play in generating organismal phenotypes is difficult to know. Indeed, only a subset of the downstream molecular consequences of a quantitative trait polymorphism may mediate its effects. Thus, even when changes in gene expression can be attributed to a single quantitative trait polymorphism, correlative and causative changes must be distinguished to fully understand the molecular basis of a trait.

In yeast, hydrogen sulfide (H<sub>2</sub>S) production varies among strains and has a negative impact on wine quality (Jiranek *et al.*, 1995; Spiropoulos & Bisson, 2000). We previously found that a strain isolated from a vineyard in Italy, M22, expresses genes in the sulfur assimilation pathway at high levels and produces high levels of H<sub>2</sub>S, a metabolite produced by the sulfur assimilation pathway (Fay *et al.*, 2004). H<sub>2</sub>S-producing strains have been shown to produce dark-brown colonies when grown in the presence of lead acetate (Ono *et al.*, 1991) or on BiGGY medium, which uses bismuth as an indicator of sulfide production (Jiranek *et al.*, 1995). Similarly, M22 produces reddish-brown or rust-colored colonies in the presence of copper sulfate. A previous work has shown that a wild-type laboratory strain produces brown-colored colonies when grown on copper-containing synthetic complete medium due to copper-sulfur mineral complexes on the cell surface (Yu *et al.*, 1996). These studies support the hypothesis that rust coloration is due to overproduction of H<sub>2</sub>S, leading to the precipitation of H<sub>2</sub>S and copper ions (Fay *et al.*, 2004). However, the relationship between expression of the sulfur assimilation genes, H<sub>2</sub>S production and colony color depends on the environment and strain background (Spiropoulos & Bisson, 2000; Linderholm *et al.*, 2008).

To dissect the relationship between changes in gene expression and the production of rust-colored colonies, we conducted a genetic analysis using M22. Based on a cross between M22 and two strains that are white-colored in the presence of copper sulfate, we found rust coloration cosegregates with another M22 phenotype, a drug-dependent delay in growth that occurs across a variety of pharmacological compounds (Kim & Fay, 2007). Quantitative trait mapping, complementation, and a single-nucleotide allele

replacement showed that an amino acid polymorphism in cystathione β-synthase (*CYS4*), I123N, causes both drug sensitivity (Kim & Fay, 2007) and rust-colored colonies (Supporting Information, Fig. S1).

*CYS4* encodes the first step in the cysteine biosynthesis pathway, which generates cysteine from homocysteine. (See Fig. 3 for an overview of the pathway.) Cysteine regulates degradation of Met4 (Menant *et al.*, 2006), a transcriptional activator of genes in the sulfur assimilation pathway. The sulfur assimilation pathway incorporates sulfur from sulfate into homocysteine via H<sub>2</sub>S (Thomas & Surdin-Kerjan, 1997). Thus, a deficiency in cysteine provides a mechanism by which the *CYS4* polymorphism could cause feedback upregulation of the sulfur assimilation pathway and high levels of H<sub>2</sub>S production. If rust coloration is caused by a reaction between copper and H<sub>2</sub>S ions, rust coloration may be mediated by upregulation of the sulfur assimilation pathway. However, the mechanism by which rust color is produced could be more complex; both cysteine and glutathione play critical roles in monitoring and regulating the oxidative/reductive state of a cell (Toledano *et al.*, 2007); hence, deficiencies in cysteine and glutathione could result in an alteration of other cellular processes that are required for the production of rust-colored colonies.

Here, we describe changes in gene expression caused by a nonsynonymous polymorphism in *CYS4* and show that the majority of differentially expressed genes are not required for the production of rust color. We also show that while genes in the sulfur assimilation pathway are required for the rust coloration phenotype, they are not required for the drug-sensitivity phenotype, indicating that the sulfur assimilation pathway plays a role in mediating the pleiotropic effects of the *CYS4* polymorphism.

## Materials and methods

### Strains and media

Rich medium [2% (w/v) yeast extract, 1% (w/v) peptone, and 2% (w/v) dextrose, with or without 2% (w/v) agar] was supplemented with copper sulfate (1 mM) and propargylglycine (320 μM) to generate propargylglycine medium for the colony color assays. For chemical complementation, L-cysteine and L-glutathione (Sigma) were dissolved in distilled water, filter-sterilized, and added to rich medium. M22 is a homothallic diploid isolated from a vineyard in Italy (Fay & Benavides, 2005). An allele replacement strain (M22-*CYS4::NI23I*) was generated by directly transforming M22-*CYS4Δ* by fusion PCR fragments containing the desired *CYS4* allele and by selecting on complete media followed by color selection (Kim & Fay, 2007). Deletion of *CYS4*, *MET10*, *MET2*, and *MET17* in M22 was induced

using the *kanMX* deletion cassette and selected on the rich medium supplemented with G418 (Wach *et al.*, 1994).

### Expression analysis

Overnight cultures of M22, M22-*CYS4* $\Delta$ , and M22-*CYS4::N123I* were diluted in rich medium to 10% (v/v) and grown for 3 h at 30 °C. Total RNA was isolated using an RNeasy mini kit (Qiagen), reverse transcribed, labeled with Cy3 or Cy5 fluorescent dyes (Genisphere), and hybridized to epoxy-coated slides (MWG Biotech) spotted with 6388, 70mer oligos (Qiagen-Operon). Arrays were scanned using a ScanArray Express laser scanner (Perkin Elmer). Microarrays were manufactured and hybridizations were completed in the microarray core facility in Washington University's Genome Center. A dye-swap was performed for one of the three replicates of each strain, where Cy3 was used instead of Cy5 to label the reference RNA, a pool of RNA from all samples. Individual array was normalized using the LOWESS algorithm to control for intensity-dependent dye effect (Dudoit *et al.*, 2002). The log-transformed median ratios of normalized two-channel intensities were used for further analysis. Significant differences in gene expression among strains (M22, M22-*CYS4* $\Delta$ , and M22-*CYS4::N123I*) were estimated using ANOVA with the following model:  $y_i = \mu + V_i + \varepsilon_i$ , where  $y_i$  is the ratio of transcripts in strain  $i$  compared with the reference pool,  $\mu$  is the average ratio across all strains,  $V_i$  is the effect of strain  $i$  on the transcript ratio, and  $\varepsilon_i$  is the error. Significant variation in gene expression among the strains was identified using the false discovery rate (FDR) method of Benjamini–Hochberg (Benjamini & Hochberg, 1995). Tukey's *post hoc* honest significant difference method implemented in R (<http://cran.r-project.org/>) was used to identify which of the three pairs of strains were different from one another. Gene expression data have been deposited in Gene Expression Omnibus (GEO, <http://www.ncbi.nlm.nih.gov/geo>) and are accessible through the GEO Series accession number GSE14102.

### Enhancer/suppressor screen

The 4664 homozygous *kanMX*-deletion strains (BY4743, *MATa* $\alpha$ , *his3* $\Delta$ 1/*his3* $\Delta$ 1, *leu2* $\Delta$ 0/*leu2* $\Delta$ 0, *lys2* $\Delta$ 0/*LYS2*, *MET15/met15* $\Delta$ 0, *ura3* $\Delta$ 0/*ura3* $\Delta$ 0) were grown on rich media plates (384/plate) for 24 h at 30 °C. Colonies were replica-plated to propargylglycine medium using a RoToR robot (Singer Instruments, UK). Photographs were taken after 24-h growth, images were cropped, and red and green channels were saved separately using PHOTOSHOP CS2 (Adobe Systems Inc.). SPOTFINDER (Saeed *et al.*, 2003) was used to grid the colonies and measure color intensities. Rust coloration of each strain was measured using the green channel intensities

divided by size of the colony. To control for edge effects from the plate, each colony was normalized by either the median of the row or column of the 384 strains in each plate, depending on which deviated more from the overall background. Enhancers and suppressors were identified by those strains showing a green intensity of  $< 0.80$  and  $> 1.20$ , respectively, from photographs of the strains grown on propargylglycine medium. At these cutoffs, the deletion strains showed slight but noticeably darker or lighter levels of rust color. At a cutoff of 0.60 and 1.30, the strains showed large, easily recognized effects on rust coloration. To assess the significance of deletions with slight effects on rust coloration, a subset of 1494 strains were phenotyped in replicate. The replicates showed an average SE of 0.012 and there were no cases where a single green intensity measurement showed a deviation  $> 0.20$  from the mean of the five replicates. The probability of observing a single value  $< 0.80$  or  $> 1.20$  from permutations of the error, as measured by each strain's deviation from the genotype mean, was  $2.7 \times 10^{-4}$  and  $2.6 \times 10^{-4}$ , respectively.


### Gene ontology analysis

Significantly enriched classes of genes were identified using gene ontology terms, and *P*-values from the hypergeometric distribution followed by the Bonferroni correction as implemented in the *Saccharomyces* Genome Database (<http://www.yeastgenome.org/>). For the analysis of enhancers and suppressors, named genes in the homozygous deletion collection were used as the background set by which significantly enriched gene ontology terms were identified.

## Results

### A deficiency in cysteine biosynthesis causes rust coloration in M22

To characterize the molecular mechanisms by which a single amino acid polymorphism within cysteine  $\beta$ -synthase (*Cys4*) causes rust coloration, we compared two strains that are identical except for the causative nonsynonymous site within *CYS4*. We previously showed that the *CYS4* amino acid polymorphism causes a deficiency in glutathione, which can be synthesized from cysteine, and that supplementation with either cysteine or glutathione eliminates drug sensitivity (Kim & Fay, 2007). Similarly, both cysteine and glutathione suppress rust coloration in M22 and strains lacking *CYS4* (Fig. 1). Because glutathione can be converted back into cysteine (Ganguli *et al.*, 2007), these results imply that rust coloration is mediated by a deficiency in intracellular levels of either glutathione or cysteine.



**Fig. 1.** Cysteine and glutathione suppress rust coloration. Deletion of *CYS4* in M22, S288c, and YPS163 results in rust coloration, similar to M22, on rich medium supplemented with 1 mM copper sulfate. The level of rust coloration is reduced in medium supplemented with 8 mM L-cysteine or 2 mM L-glutathione. Rust coloration is difficult to see in YPS163, which shows reduced growth in the presence of copper sulfate.

### Differential gene expression caused by a quantitative trait polymorphism

To characterize the molecular consequences of the *CYS4* amino acid polymorphism, we measured genome-wide changes in gene expression levels. We compared expression profiles of M22 (rust colored) to an allele replacement strain (M22-*CYS4*::*N123I*, white colored). For reference, we also compared the expression profile of the M22 allele-replacement strain to M22 with *CYS4* deleted (M22-*CYS4*Δ, rust colored). From three replicate comparisons of M22 with M22-*CYS4*::*N123I*, we identified 24 genes differentially expressed at an FDR of 5% and 141 genes at an FDR of 20% (Table S1). Compared with the white-colored allele replacement, deletion of *CYS4* caused a larger set of 45 and 189 changes in gene expression at a 5% and 20% FDR, respectively (Table S1). The average fold change for the 24 and 45 differentially expressed genes were 2.1 and 2.3, respectively. The intersection between the two sets of genes is large; 21 genes intersect at a 5% FDR and 75 genes intersect at a 20% FDR. The larger number of expression differences between M22-*CYS4*Δ and the allele replacement as well as the large and directionally consistent overlap (71/75) with these expression differences between M22 and the allele replacement is consistent with the M22 allele of *CYS4* being a hypomorph, as suggested by previous observations based on glutathione levels (Kim & Fay, 2007).

The 75 genes that differ between the two rust-colored strains and the white allele-replacement strain are significantly enriched for genes involved in sulfur metabolic

processes ( $P = 1.3 \times 10^{-4}$ ) and siderophore-iron transport ( $P = 0.0058$ , see Materials and methods). Genes that function in sulfur assimilation and metabolism of sulfur-containing amino acids, methionine and cysteine, are upregulated in M22 and M22-*CYS4*Δ (Table 1), consistent with cysteine-dependent feedback regulation by Met4 (Meinant *et al.*, 2006). Genes that function in siderophore-iron transport are likely upregulated in response to glutathione deficiency because glutathione is required for the maturation of cytoplasmic iron-sulfur-containing proteins, and glutathione deficiency causes activation of iron transporters by Aft1 and Aft2 (Sipos *et al.*, 2002; Rutherford *et al.*, 2005).

### Identification of enhancers and suppressors of rust coloration

To identify genes that affect rust coloration, we screened the yeast deletion collection (Winzeler *et al.*, 1999) for enhancers and suppressors of the rust coloration phenotype. Because the deletion collection is white in the presence of copper sulfate, we used propargylglycine, which inhibits cystathionine  $\gamma$ -lyase (*CYS3*) and cysteine biosynthesis, to phenocopy the M22 allele of *CYS4*. Figure S2 shows that similar to a deficiency in *CYS4*, propargylglycine causes decreased levels of glutathione and makes it possible to identify deletions that enhance or suppress rust coloration.

Out of 4664 homozygous diploid deletion strains, we identified 120 enhancers and 304 suppressors of rust coloration (Fig. 2, Table S2). Of the 98 enhancers that are named genes, 23 are involved in protein catabolism through endosomal or vacuolar functions, and the most significant gene ontology enrichment is ubiquitin-dependent protein catabolic processes via the multivesicular body pathway ( $P = 5.8 \times 10^{-7}$ , Table 2). However, the four strongest enhancers are all genes involved in the production of cysteine. *CYS3* and *CYS4* generate cysteine from homocysteine, *DUG2* generates cysteine by degradation of glutathione (Ganguli *et al.*, 2007), and *PDX3* produces pyridoxal 5'-phosphate (Vitamin B6), a prosthetic group required for Cys4 function (McClellan *et al.*, 2000). The list of enhancers also includes a number of genes that function in methionine or cysteine biosynthesis (Table 2), and is enriched for genes involved in sulfur amino acid metabolic processes ( $P = 0.003$ ). The enhancement of rust coloration by deletion of genes involved in cysteine biosynthesis as well as two genes involved in glutathione degradation, *DUG2* and *DUG3*, suggest that rust coloration results from low levels of cysteine rather than glutathione (Fig. 3).

Suppressors of rust coloration include 267 named genes, the majority of which are related to mitochondrial functions (Table 2). Many of the mitochondrial-related genes function in the ribosome (29 genes), cytochrome complexes (12 genes), translation (six genes), and ATP synthase (10 genes).

**Table 1.** Rust coloration and drug sensitivity are affected by different sets of *CYS4*-induced genes

Gene class	Gene name	Gene expression relative to M22- <i>CYS4::N123I</i>		Gene deletion phenotype	
		M22	M22- <i>CYS4Δ</i>	Rust color	Drug sensitivity
Sulfate transport	<i>SUL1</i>	4.1	16.7*		Resistant
	<i>SUL2</i>	1.9	3.1*		
	<i>OAC1</i>	1.4*	1.1		
Sulfate assimilation	<i>MET3</i>	7.1*	3.3	WW	Resistant
	<i>ECM17</i>	6.0 <sup>†</sup>	5.1 <sup>†</sup>	WW	
	<i>MET10</i>	4.5 <sup>†</sup>	4.5 <sup>†</sup>	WW	
	<i>MET22</i>	3.1*	2.0	W	
	<i>MET16</i>	2.4*	1.9*	W	
Homocysteine biosynthesis	<i>MET17</i>	4.6 <sup>†</sup>	2.5 <sup>†</sup>	R	Resistant
	<i>IRC7</i>	1.8 <sup>†</sup>	1.4 <sup>†</sup>		
	<i>MET2</i>	1.6	2.1 <sup>†</sup>		
Methionine biosynthesis	<i>MET6</i>	5.1*	2.0		
S-Adenosylmethionine biosynthesis	<i>SAM1</i>	1.7*	1.2	RR	
	<i>SAM2</i>	1.8*	2.0*		Resistant
Methionine salvage	<i>ADI1</i>	1.7*	1.0		
Cysteine biosynthesis	<i>CYS4</i>	2.5 <sup>†</sup>	0.05 <sup>†</sup>	RR	Sensitive
	<i>CYS3</i>	1.5*	1.6*	RR	Sensitive
Glutathione biosynthesis	<i>GSH1</i>	1.6	2.1*		Sensitive
Siderophore-iron transport	<i>ARN1</i>	1.5 <sup>†</sup>	3.1 <sup>†</sup>		Resistant
	<i>SIT1</i>	2.9 <sup>†</sup>	2.7 <sup>†</sup>		
	<i>ENB1</i>	2.5*	2.4*		
	<i>FIT2</i>	1.6	8.5 <sup>†</sup>		
	<i>FIT3</i>	1.4	7.8 <sup>†</sup>		

\*Genes significant at 20% FDR.

<sup>†</sup>Genes significant at 5% FDR.

Gene expression ratios are the average of three replicates and the mean squared error is 0.13, averaged across genes. Rust coloration and drug sensitivity phenotypes were measured using strains from the homozygous diploid deletion collection. RR indicates a strong enhancer ( $< 0.6$ ) and WW a strong suppressor ( $> 1.3$ ), as measured by the intensity of the green channel from photographs. R indicates a moderate enhancer ( $< 0.8$ ) and W a moderate suppressor ( $> 1.2$ ). Drug sensitivity was measured by the growth delay induced by 2 mM atenolol supplemented to rich medium. Strains with  $< 4$  h of growth delay are classified as resistant and those with  $> 9$  h of growth delay are classified as sensitive.

Although there is enrichment for genes involved in cellular respiration ( $P = 2.0 \times 10^{-15}$ ) and genes that function in the mitochondria ( $P = 1.1 \times 10^{-35}$ ), there are many genes required for respiration that did not suppress rust coloration. Thus, the suppression of rust color is not simply due to the reduced rate of growth of petite mutants. In addition to mitochondrial-related genes, there is also enrichment for genes involved in gene expression ( $P = 7.2 \times 10^{-8}$ ) and sulfate assimilation ( $P = 3.0 \times 10^{-5}$ ). All six genes that function in the assimilation of sulfur from sulfate into homocysteine are suppressors of rust coloration (Table 2, Fig. 3). The last step in the sulfur assimilation pathway is the formation of homocysteine from  $H_2S$  and *O*-acetyl-L-homoserine by Met17, which is an enhancer of rust coloration. The suppression of rust coloration by deletion of genes that lead to the production of  $H_2S$  combined with the enhancement of rust coloration by deletion of *MET17* indicates that  $H_2S$  is a key metabolite involved in the production of rust coloration (Fig. 3).

### Differentially expressed genes necessary for the production of rust coloration

If genes that are differentially expressed as a result of the *CYS4* polymorphism play a role in generating the rust coloration phenotype, then deletion of these genes should either enhance or suppress rust coloration. Seven of the 120 enhancers and eight of the 304 suppressors were differentially expressed between both the rust- (M22, M22-*CYS4Δ*) and the white-colored (M22-*CYS4::N123I*) strains. Genes that are upregulated in M22 should suppress rust coloration when deleted if their expression levels are involved in the production of rust color. Of the eight suppressors, seven are expressed at higher levels in the rust-colored strains and five of these function in the conversion of sulfate into  $H_2S$  before assimilation into homocysteine (Fig. 3). The other two suppressors are *TUF1* and *YTA12*, required for translation of mitochondrial proteins and assembly of mitochondrial enzyme complexes, respectively. Only one of the enhancers,

*SWI3*, was expressed at lower levels in the two rust-colored strains.

The identification of differentially expressed genes necessary for the production of rust coloration provides insight into the mechanism by which the *CYS4* polymorphism results in rust coloration. The upregulation of both enhancers and suppressors in the same biochemical pathway suggests that  $H_2S$  is involved in the production of rust coloration. To confirm that  $H_2S$  production is required for rust coloration in M22, we deleted *MET2*, *MET17*, and *MET10* in M22. The suppression of rust coloration by *MET10* and enhancement of rust coloration by *MET2* or *MET17* shows that rust coloration in M22 depends on  $H_2S$  production (Fig. 4). To confirm the requirement for mito-


chondrial function in M22 and test whether mitochondrial function is required for  $H_2S$  production, we deleted both *TUF1* and *YTA12* in M22. Surprisingly, neither deletion suppressed rust coloration. Rust coloration of M22 was also unaffected by deletion of two other genes, *MRPL38* and *MRPS17*, that are required for rust coloration in the deletion collection background.

### Pleiotropy at the molecular level

In addition to rust coloration, the M22 allele of *CYS4* also causes a delay in growth upon exposure to a variety of different pharmacological compounds (Kim & Fay, 2007). While both the rust coloration and drug-sensitivity phenotypes are caused by a deficiency in cysteine/glutathione, the molecular basis of these two traits may not be identical. To determine whether any of the differentially expressed genes that function in sulfur assimilation or iron transport affect drug sensitivity, we measured growth in the presence and the absence of atenolol, a  $\beta$ -adrenergic receptor antagonist that causes a delay in growth dependent on the *CYS4* amino acid polymorphism (Kim & Fay, 2007). Four of the genes involved in the production of sulfur-containing amino acids and two of the genes involved in iron transport increase drug resistance when deleted. However, most of the genes in the sulfur assimilation pathway that affect rust coloration do not affect drug sensitivity (Table 1). In contrast, deletions of genes required for glutathione biosynthesis, *CYS3*, *CYS4* and *GSH1*, increase drug sensitivity. Thus, while both rust coloration and drug sensitivity result from a deficiency in cysteine, they each have a distinct molecular basis because they are enhanced and suppressed by different sets of genes.

### Discussion


Dissecting the molecular mechanisms by which a single quantitative trait polymorphism influences a trait of interest provides insight into those biochemical or signaling pathways that underlie a trait. We identified a complex set of downstream changes in gene expression caused by a pleiotropic quantitative trait polymorphism. We then showed that a subset of the differentially expressed genes in the same




**Fig. 2.** Phenotypic distribution of enhancers and suppressors of rust coloration in the homozygous yeast deletion collection. Rust coloration was measured by the average intensity of the green channel from RGB photographs of a strain divided by the median of all strains in the same row or column (see Materials and methods for details). Enhancers are shown in red and suppressors in blue. Photographs of selected strains show five replicates per strain, illustrating the mean and SD of the rust coloration phenotype.

**Table 2.** Functional classification of enhancers and suppressors

Pathway/function	Subclass	Enhancers	Suppressors
Sulfate assimilation and sulfur metabolism	Enzyme	<i>SAM1</i> , <i>CYS3/4</i> , <i>DUG2/3</i> , <i>MET17</i> , <i>GLO2</i>	<i>MET3/10/14/16/22</i> , <i>ECM17</i>
	Regulation	<i>BUL1</i> , <i>STP1</i>	<i>MET28/32</i>
	Siroheme biosynthesis		<i>MET1/8</i>
	Purine and adenosine biosynthesis	<i>ADE1</i>	<i>ADE3/5/6</i>
	Transport	<i>MUP1</i>	
	SAM-dependent methyl transfer	<i>CHO2</i>	
Vacuolar, golgi, protein sorting		23 genes	
Mitochondrial			221 genes
Total		120 genes	304 genes



**Fig. 3.** Genes involved in sulfur assimilation and biosynthesis of cysteine and glutathione. Asterisks and crosses indicate suppressors and enhancers of rust color, respectively. Genes not included in the screen are shown in parentheses. Genes upregulated in M22 or in M22-CYS4 $\Delta$  are indicated by a triangle. Dotted lines show ubiquitin-mediated negative regulation of Met4 by cysteine and positive regulation of the sulfur assimilation pathway by the Met4 complex. APS, adenosine 5'-phosphosulfate; PAPS, 3'-phosphoadenosine-5'-phosphosulfate; SAM, S-adenosylmethionine;  $\gamma$ -GluCys,  $\gamma$ -glutamylcysteine; SAH, S-adenosylhomocysteine.



**Fig. 4.** Deletion of *MET2* or *MET17* enhances rust coloration, and deletion of *MET10* suppresses rust coloration in M22. The four meiotic progeny of each heterozygous M22 deletion strain were plated on rich medium with 1 mM copper sulfate and labels show strain genotypes.

biochemical pathway is required for the production of rust coloration but not drug sensitivity. These results provide insight into the molecular basis of pleiotropy.

### The molecular basis of rust coloration

The suppression of rust coloration by deletion of genes required for H<sub>2</sub>S production and enhancement of rust coloration by deletion of genes involved in utilization of H<sub>2</sub>S implies that rust coloration is mediated by H<sub>2</sub>S production. Copper-dependent changes in yeast colony color are thought to be due to copper-sulfur mineralization at the cell surface (Ashida *et al.*, 1963; Yu *et al.*, 1996). Consistent with

this possibility, electron micrographs show dark granules at the cell surface in M22 but not the white-colored M22 allele replacement strain (Fig. S3). However, H<sub>2</sub>S production may not be sufficient for the production of rust color. The rust-colored compound may also depend on other cellular perturbations such as a deficiency in glutathione, a potent antioxidant, or altered levels of metabolites in the S-adenosylmethionine pathway (Christopher *et al.*, 2002). In the yeast deletion collection background, rust coloration is dependent on some aspect of mitochondrial function, but not in M22. The large number of suppressors with mitochondrial functions makes it difficult to know which mitochondrial function is required for rust coloration. One possibility is that CuS mineralization occurs in the mitochondrion or requires iron-sulfur-containing proteins synthesized in mitochondria (Kispal *et al.*, 1999). However, it is also possible that mitochondrial mutants suppress rust coloration by affecting H<sub>2</sub>S production, either by reducing the need for glutathione or by affecting the sulfur assimilation genes through mitochondrial retrograde regulation (Liu & Butow, 2006).

In M22, H<sub>2</sub>S production may be elevated due to differential regulation of the sulfur assimilation pathway. Alternatively, the M22 allele of *CYS4* may cause homocysteine utilization to be the rate-limiting step in sulfur assimilation.

In this scenario, upregulation of the sulfur assimilation genes would have little or no effect on H<sub>2</sub>S levels. Consistent with this latter possibility, upregulation of the sulfur assimilation pathway in another Italian wine strain due to a frameshift in the extracellular amino acid sensor, *SSY1* (Brown *et al.*, 2008), does not lead to rust coloration or noticeable levels of H<sub>2</sub>S production (K. Brown, pers. commun.).

Our enhancer and suppressor screen implies that hundreds of genes with a wide range of biological functions affect rust coloration. In many cases, enhancement and suppression of rust coloration may be mediated by changes in cysteine/glutathione homeostasis and H<sub>2</sub>S production. Evidence for this possibility can be found in the overlap between enhancers and suppressors of rust coloration and two other screens of the yeast deletion collection. A screen for genes involved in glutathione homeostasis identified 276 genes that excrete high levels of glutathione, and one of the largest functionally related groups contains genes involved in the secretory pathway or vacuolar protein sorting (Perone *et al.*, 2005), similar to enhancers of rust coloration. Of the 276 genes, 96 overlap with enhancers or suppressors of rust coloration. Another screen for genes involved in H<sub>2</sub>S production identified 89 genes, many of which function in the biosynthesis of sulfur-containing amino acids (Linderholm *et al.*, 2008). Of the 89 deletions that cause high levels of sulfite reductase activity, 15 overlap with enhancers of rust coloration. These comparisons imply that some but not all of the enhancers and suppressors of rust coloration affect cysteine/glutathione levels and H<sub>2</sub>S production in the absence of propargylglycine-induced inhibition of cysteine biosynthesis.

### The role of gene expression in pleiotropy

We found a large number of genes that are differentially expressed as a consequence of the *CYS4* polymorphism. The upregulation of genes that suppress rust coloration when deleted implies that the differential expression of genes in the sulfur assimilation pathway plays an important role in the production of rust color. However, other *CYS4*-induced changes in gene expression may have no downstream phenotypic consequences. For example, the upregulation of genes that enhance drug sensitivity when deleted, for example *CYS3* or *GSH1*, suggests that some expression changes are a response to alterations in cysteine homeostasis but are not responsible for drug sensitivity. Finally, some *CYS4*-induced changes in gene expression may only have phenotypic consequences under certain environmental conditions; for example rust coloration is only observed in the presence of copper sulfate, but upregulation of the sulfur assimilation genes occurs both in the presence and the absence of copper sulfate (Fay *et al.*, 2004). Thus, the

differential expression of siderophore-iron transporters may also produce a downstream phenotype when measured under the right condition. The upregulation of *SIT1* and *FIT2* combined with the higher level of drug-resistance when deleted implies that the differential expression of these two genes may be involved in drug sensitivity.

### Implications for phenotypic variation in other yeast isolates

What is the relevance of our results to phenotypic variation in other strains of yeast? While the rust coloration phenotype is rare, progeny of M22 crossed with two white-colored strains show a continuous distribution of rust coloration phenotypes in those strains that inherit the M22 allele of *CYS4* (Fig. S4). Furthermore, natural yeast isolates show significant variation in levels of intracellular glutathione (Fig. S5), which affects metabolism of a wide range of pharmacological compounds (Kim & Fay, 2007), and H<sub>2</sub>S, which affects wine production (Mendes-Ferreira *et al.*, 2002). Thus, our results provide insight into the genes, pathways, and processes that affect population variation in cysteine/glutathione metabolism and H<sub>2</sub>S production. However, we also found evidence for significant effects of genetic background. Deletions of four mitochondrial genes suppress rust coloration in BY4743, the strain used to make the deletion collection, but do not suppress rust coloration in M22. Thus, differences in genetic background may have resulted in missed M22-specific enhancers or suppressors, and presents a significant challenge to obtaining a comprehensive understanding of the molecular basis of cysteine-related phenotypes.

How common are pleiotropic, background-dependent quantitative trait nucleotides? A nonsynonymous polymorphism in *MKT1* shares many of the characteristics of the *CYS4* polymorphism: extensive pleiotropy, background-dependent effects, and numerous changes in gene expression levels (Deutschbauer & Davis, 2005; Sinha *et al.*, 2006; Demogines *et al.*, 2008; Smith & Kruglyak, 2008). However, small-effect polymorphisms may have quite different characteristics. For example, quantitative trait polymorphisms with small effects show less pleiotropy than those of large effect (Wagner *et al.*, 2008).

### Acknowledgements

We thank M. Johnston for sharing the yeast deletion collection, B. Cohen for sharing microarrays, L. Riles and J. Dover for technical support, W. Beatty in the Molecular Microbiology Imaging Facility for electron microscopy service, C. Hittinger, M. Dorsett, D. Swain, and B. Engle (all at Washington University) for critical reading and helpful discussions, and K. Brown (Harvard University) for

examining phenotypes in another Italian wine strain. This work was supported by a National Institute of Health grant (80669) to J.C.F.

## References

- Ashida J, Higashi N & Kikuchi T (1963) An electronmicroscopic study on copper precipitation by copper-resistant yeast cells. *Protoplasma* **57**: 27–32.
- Benjamini Y & Hochberg Y (1995) Controlling the false discovery rate – a practical and powerful approach to multiple testing. *J Roy Stat Soc B* **57**: 289–300.
- Birrell GW, Brown JA, Wu HI, Giaever G, Chu AM, Davis RW & Brown JM (2002) Transcriptional response of *Saccharomyces cerevisiae* to DNA-damaging agents does not identify the genes that protect against these agents. *P Natl Acad Sci USA* **99**: 8778–8783.
- Brown KM, Landry CR, Hartl DL & Cavalieri D (2008) Cascading transcriptional effects of a naturally occurring frameshift mutation in *Saccharomyces cerevisiae*. *Mol Ecol* **17**: 2985–2997.
- Carroll AS, Bishop AC, DeRisi JL, Shokat KM & O’Shea EK (2001) Chemical inhibition of the Pho85 cyclin-dependent kinase reveals a role in the environmental stress response. *P Natl Acad Sci USA* **98**: 12578–12583.
- Cavalieri D, Townsend JP & Hartl DL (2000) Manifold anomalies in gene expression in a vineyard isolate of *Saccharomyces cerevisiae* revealed by DNA microarray analysis. *P Natl Acad Sci USA* **97**: 12369–12374.
- Christopher SA, Melnyk S, James SJ & Kruger WD (2002) S-adenosylhomocysteine, but not homocysteine, is toxic to yeast lacking cystathionine beta-synthase. *Mol Genet Metab* **75**: 335–343.
- de Haan G, Bystrykh LV, Weersing E *et al.* (2002) A genetic and genomic analysis identifies a cluster of genes associated with hematopoietic cell turnover. *Blood* **100**: 2056–2062.
- Demogines A, Smith E, Kruglyak L & Alani E (2008) Identification and dissection of a complex DNA repair sensitivity phenotype in Baker’s yeast. *PLoS Genet* **4**: e1000123.
- Deutschbauer AM & Davis RW (2005) Quantitative trait loci mapped to single-nucleotide resolution in yeast. *Nat Genet* **37**: 1333–1340.
- Dudoit S, Yang YH, Callow M & Speed T (2002) Statistical methods for identifying differentially expressed genes in replicated cDNA microarray experiments. *Stat Sinica* **12**: 111–139.
- Fay JC & Benavides JA (2005) Evidence for domesticated and wild populations of *Saccharomyces cerevisiae*. *PLoS Genet* **1**: 66–71.
- Fay JC, McCullough HL, Sniegowski PD & Eisen MB (2004) Population genetic variation in gene expression is associated with phenotypic variation in *Saccharomyces cerevisiae*. *Genome Biol* **5**: R26.
- Ganguli D, Kumar C & Bachhawat AK (2007) The alternative pathway of glutathione degradation is mediated by a novel protein complex involving three new genes in *Saccharomyces cerevisiae*. *Genetics* **175**: 1137–1151.
- Giaever G, Chu AM, Ni L *et al.* (2002) Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* **418**: 387–391.
- Jiraneek V, Langridge P & Henschke PA (1995) Regulation of hydrogen sulfide liberation in wine-producing *Saccharomyces cerevisiae* strains by assimilable nitrogen. *Appl Environ Microb* **61**: 461–467.
- Kim HS & Fay JC (2007) Genetic variation in the cysteine biosynthesis pathway causes sensitivity to pharmacological compounds. *P Natl Acad Sci USA* **104**: 19387–19391.
- Kispal G, Csere P, Prohl C & Lill R (1999) The mitochondrial proteins Atm1p and Nfs1p are essential for biogenesis of cytosolic Fe/S proteins. *EMBO J* **18**: 3981–3989.
- Lee SI, Dudley AM, Drubin D, Silver PA, Krogan NJ, Pe’er D & Koller D (2009) Learning a prior on regulatory potential from eQTL data. *PLoS Genet* **5**: e1000358.
- Linderholm AL, Findleton CL, Kumar G, Hong Y & Bisson LF (2008) Identification of genes affecting hydrogen sulfide formation in *Saccharomyces cerevisiae*. *Appl Environ Microb* **74**: 1418–1427.
- Liu Z & Butow RA (2006) Mitochondrial retrograde signaling. *Annu Rev Genet* **40**: 159–185.
- McClellan KN, Janosik M, Oliveriusová J, Kery V & Kraus JP (2000) Transsulfuration in *Saccharomyces cerevisiae* is not dependent on heme: purification and characterization of recombinant yeast cystathionine beta-synthase. *J Inorg Biochem* **81**: 161–171.
- Menant A, Baudouin-Cornu P, Peyraud C, Tyers M & Thomas D (2006) Determinants of the ubiquitin-mediated degradation of the Met4 transcription factor. *J Biol Chem* **281**: 11744–11754.
- Mendes-Ferreira A, Mendes-Faia A & Leao C (2002) Survey of hydrogen sulphide production by wine yeasts. *J Food Protect* **65**: 1033–1037.
- Ono B, Ishii N, Fujino S & Aoyama I (1991) Role of hydrosulfide ions (HS<sup>-</sup>) in methylmercury resistance in *Saccharomyces cerevisiae*. *Appl Environ Microb* **57**: 3183–3186.
- Perrone GG, Grant CM & Dawes IW (2005) Genetic and environmental factors influencing glutathione homeostasis in *Saccharomyces cerevisiae*. *Mol Biol Cell* **16**: 218–230.
- Rutherford JC, Ojeda L, Balk J, Muhlenhoff U, Lill R & Winge DR (2005) Activation of the iron regulon by the yeast Aft1/Aft2 transcription factors depends on mitochondrial but not cytosolic iron–sulfur protein biogenesis. *J Biol Chem* **280**: 10135–10140.
- Saeed AI, Sharov V, White J *et al.* (2003) TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* **34**: 374–378.
- Schadt EE, Monks SA, Drake TA *et al.* (2003) Genetics of gene expression surveyed in maize, mouse and man. *Nature* **422**: 297–302.
- Sinha H, Nicholson BP, Steinmetz LM & McCusker JH (2006) Complex genetic interactions in a quantitative trait locus. *PLoS Genet* **2**: e13.

- Sipos K, Lange H, Fekete Z, Ullmann P, Lill R & Kispal G (2002) Maturation of cytosolic iron–sulfur proteins requires glutathione. *J Biol Chem* **277**: 26944–26949.
- Smith EN & Kruglyak L (2008) Gene–environmental interaction in yeast gene expression. *PLoS Biol* **6**: e83.
- Spiropoulos A & Bisson LF (2000) *MET17* and hydrogen sulfide formation in *Saccharomyces cerevisiae*. *Appl Environ Microb* **66**: 4421–4426.
- Thomas D & Surdin-Kerjan Y (1997) Metabolism of sulfur amino acids in *Saccharomyces cerevisiae*. *Microbiol Mol Biol R* **61**: 503–532.
- Toledano MB, Kumar C, Le Moan N, Spector D & Tacnet F (2007) The system biology of thiol redox system in *Escherichia coli* and yeast: differential functions in oxidative stress, iron metabolism and DNA synthesis. *FEBS Lett* **581**: 3598–3607.
- Wach A, Brachat A, Pohlmann R & Philippsen P (1994) New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast* **10**: 1793–1808.
- Wagner GP, Kenney-Hunt JP, Pavlicev M, Peck JR, Waxman D & Cheverud JM (2008) Pleiotropic scaling of gene effects and the ‘cost of complexity’. *Nature* **452**: 470–472.
- Winzeler EA, Shoemaker DD, Astromoff A *et al.* (1999) Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science* **285**: 901–906.
- Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV & Romano LA (2003) The evolution of transcriptional regulation in eukaryotes. *Mol Biol Evol* **20**: 1377–1419.
- Yu W, Farrell RA, Stillman DJ & Winge DR (1996) Identification of *SLF1* as a new copper homeostasis gene involved in copper sulfide mineralization in *Saccharomyces cerevisiae*. *Mol Cell Biol* **16**: 2464–2472.
- Yvert G, Brem RB, Whittle J *et al.* (2003) Trans-acting regulatory variation in *Saccharomyces cerevisiae* and the role of transcription factors. *Nat Genet* **35**: 57–64.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Fig. S1.** Rust coloration is affected by a nonsynonymous polymorphism in *CYS4*.

**Fig. S2.** Propargylglycine inhibits glutathione biosynthesis and causes rust coloration.

**Fig. S3.** Electron micrographs show dark granules at the cell surface in M22 but not M22-*CYS4::N123I*.

**Fig. S4.** The phenotypic distribution of homozygous recombinant strains.

**Fig. S5.** Variation in intracellular glutathione concentration among natural yeast isolates.

**Table S1.** Gene expression differences among strains.

**Table S2.** Enhancers and suppressors of rust color.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.